



Classifying new anti-tuberculosis drugs: rationale and future perspectives



Simon Tiberi^a, Anna Scardigli^b, Rosella Centis^c, Lia D'Ambrosio^{c,d},
 Marcela Muñoz-Torrico^e, Miguel Ángel Salazar-Lezama^e, Antonio Spanevello^{f,g},
 Dina Visca^f, Alimuddin Zumla^h, Giovanni Battista Migliori^{c,*}, Jose A. Caminero Luna^{ij}

^a Division of Infection, Royal London Hospital, Barts Health NHS Trust, London, UK

^b The Global Fund to Fight Aids, Tuberculosis and Malaria, Geneva, Switzerland

^c Fondazione S. Maugeri, IRCCS, Care and Research Institute, Via Roncaccio 16, 21049, Tradate, Italy

^d Public Health Consulting Group, Lugano, Switzerland

^e Tuberculosis Clinic, National Institute of Respiratory Diseases of Mexico (INER), Mexico City, Mexico

^f Department of Clinical and Experimental Medicine, University of Insubria, Varese, Italy

^g Pneumology Unit, Fondazione Maugeri, IRCCS, Care and Research Institute, Tradate, Italy

^h Division of Infection and Immunity, University College London and NIHR Biomedical Research Centre, UCL Hospitals NHS Foundation Trust, London, UK

ⁱ Pneumology Department, University Hospital of Gran Canaria "Dr Negrin", Las Palmas de Gran Canaria, Las Palmas, Spain

^j International Union against Tuberculosis and Lung Disease, Paris, France

ARTICLE INFO

Article history:

Received 5 October 2016

Received in revised form 24 October 2016

Accepted 25 October 2016

Corresponding Editor: Eskild Petersen, Aarhus, Denmark.

Keywords:

Anti-TB drugs

MDR/XDR-TB

Fluoroquinolones

Linezolid

Bedaquiline

Delamanid ;

SUMMARY

The classification of anti-tuberculosis (TB) drugs is important as it helps the clinician to build an appropriate anti-TB regimen for multidrug-resistant (MDR) and extensively drug-resistant (XDR) TB cases that do not fulfil the criteria for the shorter MDR-TB regimen. The World Health Organization (WHO) has recently approved a revision of the classification of new anti-TB drugs based on current evidence on each drug. In the previous WHO guidelines, the choice of drugs was based on efficacy and toxicity in a step-down manner, from group 1 first-line drugs and groups 2–5 second-line drugs, to group 5 drugs with potentially limited efficacy or limited clinical evidence. In the revised WHO classification, exclusively aimed at managing drug-resistant cases, medicines are again listed in hierarchical order from group A to group D. In parallel, a possible future classification is independently proposed. The aim of this viewpoint article is to describe the evolution in WHO TB classification (taking into account an independently proposed new classification) and recent changes in WHO guidance, while commenting on the differences between them. The latest evidence on the ex-group 5 drugs is also discussed.

© 2016 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The World Health Organization (WHO) has recently updated the classification of new anti-tuberculosis (TB) drugs based on a meta-analysis and expert panel recommendations.¹ During the period between the publication of the first WHO anti-TB drug classification and the revised version, an independent proposal for a new classification was made available in the literature.² Evidence for further reclassification is lacking and will only be forthcoming with data from new randomized controlled trials

(RCTs) aimed at developing better (more effective and tolerated) regimens. However, even though a new classification is not required, discussion on possible future steps has begun,² with particular focus on some of the existing second-line anti-TB drugs.

The classification of anti-TB drugs is important as it helps the clinician to build an appropriate anti-TB regimen for multidrug-resistant (MDR) and extensively drug-resistant (XDR) TB cases that do not fulfil the criteria for the shorter MDR-TB regimen.^{3,4}

The aim of this viewpoint article is to describe the evolution in WHO TB classification (taking into account an independently proposed new classification) and recent changes in WHO guidance, while commenting on the differences between them. The latest evidence on the ex-group 5 drugs is also discussed.

* Corresponding author. Tel.: +39 0331829404.

E-mail address: giovannibattista.migliori@fsm.it (G.B. Migliori).

2. Classifications

In the previous WHO guidelines (2011), the choice of drugs was based on efficacy and toxicity in a step-down manner, from group 1 to group 5 (Table 1). Group 1 included first-line drugs and groups 2–5 included second-line drugs. Group 5 included the drugs with (at the time) potentially limited efficacy or limited clinical evidence.^{3,4}

According to the new WHO drug classification (2016), patients with rifampicin-resistant or MDR-TB require a regimen with at least five effective TB medicines during the intensive phase: pyrazinamide and four core second-line TB drugs (see Table 1), one each from group A and group B, and at least two from group C. If the minimum number of effective TB medicines cannot be composed, an agent from group D2 and other agents from D3 should be added to bring the total to five. If pyrazinamide is compromised or cannot be used, the regimen can be reinforced with a drug from group C or D (preferably D2, and if not, from D3). Agents from group D1 are added if they are considered to add benefit (e.g., high-dose isoniazid in patients without high-level isoniazid resistance).¹ The total number of TB medicines included in the regimen needs to balance the expected benefit with the risk of harm and non-adherence.

Based on recent evidence on given compounds, some drugs are likely to increase or decrease in importance in the future.²

2.1. Group 1

In accordance with drug susceptibility testing (DST), all active group 1 drugs (Table 1) should be included in the regimen, taking into consideration that isoniazid, rifampicin/rifabutin, and pyrazinamide are core drugs and ethambutol is a companion drug. Streptomycin is no longer used routinely.

High-dose isoniazid can be added to an MDR/XDR-TB regimen when the *katG* mutation is not detected by line probe assay, but should not be counted as one of the four active drugs^{1,3,4} (although recent evidence suggests the mutation confers intermediate resistance only). Pyrazinamide should always be used, as DST is unreliable; however, it should not be counted as one of the four active drugs.^{1,3,4} Rifabutin should be considered if sensitivity is proven and a favourable mutation profile exists.⁵ More specifically, if rifampicin resistance is detected with rifabutin susceptibility, rifabutin should be added to the regimen, but not counted as one of the four active drugs.⁵

As the new WHO classification is aimed at managing drug-resistant cases and not all cases, as in the previous classification, group 1 drugs lose priority. In the new WHO classification they belong, in fact, to group D1.

2.2. Group A

According to the new WHO classification,¹ group A now includes fluoroquinolones and group B includes injectable second-line drugs. Fluoroquinolones (particularly the later generation fluoroquinolones such as high-dose levofloxacin, gatifloxacin, or moxifloxacin) are core drugs, demonstrating bactericidal and sterilizing activity and a good safety profile;^{2,3,6,7} their use predicts a favourable outcome in the treatment of MDR-TB.^{7–11} They are the best agents for the treatment of MDR-TB.

2.3. Group B

The second-line injectable drugs have only bactericidal and no sterilizing activity. As their safety profile is clearly worse than that of fluoroquinolones, they remain a step below them on the ranking.^{2,4,6}

Table 1

Summary of the existing classifications of anti-tuberculosis drugs (1 and 2) and possible future changes based on recent evidence (3)

(1) WHO 2011 TB drugs classification		(2) WHO 2016 TB drugs classification		(3) Possible future evolutions	
Group 1 First-line oral anti-TB drugs	<ul style="list-style-type: none"> • Isoniazid • Rifampicin • Ethambutol • Pyrazinamide 	Group A Fluoroquinolones	<ul style="list-style-type: none"> • Levofloxacin • Moxifloxacin • Gatifloxacin 	Group A Fluoroquinolones	<ul style="list-style-type: none"> • Levofloxacin • Moxifloxacin • Gatifloxacin
Group 2 Injectable anti-TB drugs (injectable or parenteral agents)	<ul style="list-style-type: none"> • Streptomycin • Kanamycin • Amikacin • Capreomycin 	Group B Second-line injectable agents	<ul style="list-style-type: none"> • Amikacin • Capreomycin • Kanamycin (Streptomycin) 	Group B Other core second-line agents	<ul style="list-style-type: none"> • Bedaquiline • Delamanid • Ethionamide/prothionamide • Cycloserine/terizidone • Linezolid • Clofazimine • Amikacin • Capreomycin • Kanamycin • Meropenem/clavulanate
Group 3 Fluoroquinolones	<ul style="list-style-type: none"> • Levofloxacin • Moxifloxacin • Gatifloxacin • Ofloxacin 	Group C Other core second-line agents	<ul style="list-style-type: none"> • Ethionamide/prothionamide • Cycloserine/terizidone • Linezolid • Clofazimine 	Group C Second-line injectable agents	<ul style="list-style-type: none"> • Pyrazinamide • Ethambutol • High-dose isoniazid
Group 4 Oral bacteriostatic second-line anti-TB drugs	<ul style="list-style-type: none"> • Ethionamide/prothionamide • Cycloserine/terizidone • <i>p</i>-Aminosalicylic acid 	Group D Add-on agents (not core MDR-TB regimen components)	<ul style="list-style-type: none"> • Pyrazinamide • Ethambutol • High-dose isoniazid 	Group D Add-on agents (not core MDR-TB regimen components)	<ul style="list-style-type: none"> • Pyrazinamide • Ethambutol • High-dose isoniazid
Group 5 Anti-TB drugs with limited data on efficacy and long-term safety in the treatment of drug-resistant TB	<ul style="list-style-type: none"> • Linezolid • Clofazimine • Amoxicillin/clavulanate • Imipenem/cilastatin • Meropenem • High-dose isoniazid • Thioacetazone • Clarithromycin 		<ul style="list-style-type: none"> • Bedaquiline • Delamanid • <i>p</i>-Aminosalicylic acid • Imipenem–cilastatin • Meropenem • Amoxicillin–clavulanate • (Thioacetazone) 		<ul style="list-style-type: none"> • <i>p</i>-Aminosalicylic acid • Amoxicillin–clavulanate • Rifabutin

WHO, World Health Organization; MDR-TB, multidrug-resistant tuberculosis.

Furthermore, the second group in the drug hierarchy (group B) may in future include three core oral medicines, linezolid, bedaquiline, and delamanid (and eventually sutezolid, tedizolid, and pretomanid), if these drugs prove to be more effective and better tolerated than the injectables.² This has been proposed in a recent article as group 3 (similar to the current group B, because in this article group 1 is composed of first-line drugs).²

Moreover, having an oral group B could mean that it may soon be possible to have an injectable-free regimen to treat MDR-TB patients; this will mean potentially less toxicity, less monitoring, and fewer hospital stays and visits, and possibly better adherence.

2.4. Group C

Linezolid, clofazimine, ethionamide/prothionamide, and cycloserine/terizidone are included in group C. Linezolid is a core oral drug with bactericidal and sterilizing action;² there is ample evidence of its efficacy, including RCTs and meta-analyses.^{12,13} The drug is generic and active, its documented toxicity being the primary barrier to continued use. However, this can be mitigated with lower doses and therapeutic drug monitoring.^{13,14} Linezolid has the efficacy needed to be part of a future hypothetical all-oral group B. With effectiveness in mind, the second drug in this group could be ethionamide/prothionamide, which has moderate bactericidal activity but with an appreciable toxicity profile.⁴ The third agent could be clofazimine, as this has possible sterilizing activity and good tolerance. The last one could be cycloserine, the worst of the four, with practically no bactericidal or sterilizing activity and with a poor toxicity profile.^{2,4}

2.5. Group D

Group D1 includes pyrazinamide, ethambutol, and high-dose isoniazid. Group D2 includes bedaquiline and delamanid. These may have the efficacy needed to be part of a future hypothetical all-oral group B, given that evidence on their safety is growing, although it is still incomplete. Data regarding their safe combination and whole treatment duration over the recommended 6 months are gradually emerging.

Recent case reports show the safe and effective use of bedaquiline up to 18 months¹⁵ and the concomitant use of both group D2 drugs.¹⁶ Bedaquiline has all of the characteristics of a core drug, targeting both actively replicating and dormant bacilli.¹⁷ RCTs provide some evidence on efficacy and safety,^{18,19} while additional experience derived from observational studies and compassionate use programmes completes the current picture.^{15,20–22} Concerns regarding the safety of bedaquiline arose from the unexplained higher number of deaths in the bedaquiline group in a licensing study.¹⁸

The most common adverse reaction associated with bedaquiline is a QTc interval increase on electrocardiogram.^{18,19} An RCT with long-term follow-up to 120 weeks reported that bedaquiline was well tolerated and led to good outcomes.²⁰ The reporting of adverse events is crucial for recommendations on bedaquiline use;²³ therefore monitoring, active pharmacovigilance, and proper management of adverse reactions are foremost among the five criteria in place for the use of this agent. Finally, a possible issue is cross-resistance with clofazimine²⁴ (although recent evidence tends to suggest that this may not be clinically relevant) and the increased QT prolongation with clofazimine and bedaquiline combination (which appears to be a causative factor leading to increased QTc in previous bedaquiline studies).²⁵

Delamanid can also be considered a core drug because of its bactericidal and sterilizing activity; it does not show cross-resistance with other anti-TB drugs.²⁶ Some RCTs and observational studies have addressed its efficacy,²⁶ and there are also some

positive experiences from its compassionate use.²⁶ WHO recommendations on delamanid use²⁷ include the same five implementation criteria as in the case of bedaquiline.²³

As mentioned above, a significant limitation of bedaquiline and delamanid use is that, so far, they can be utilized only for the first 6 months of treatment. When treating XDR-TB patients, these drugs are added to an optimized background regimen that often includes weak and poorly tolerated medications (from the limited options remaining), with severe side effects mandating interruption of either the whole treatment or of the offending compound. When this happens, the regimen becomes even weaker and, once bedaquiline or delamanid is completed after 6 months, the regimen is prone to fail.

The possibility of maintaining bedaquiline and/or delamanid for the entire duration of treatment will be an important step forward, as well as their use in patients with MDR-TB patterns of resistance beyond XDR-TB,¹⁰ or the so-called pre-XDR-TB (e.g., TB sustained by strains resistant to either fluoroquinolones or second-line injectables).^{2,15,16}

After the proposed reclassification of groups 2 and 3 to group B and group A, respectively, the injectable second-line drugs might have the characteristics of a future group C, as recently suggested (Table 1).²

Indeed, they remain core second-line drugs thanks to their bactericidal activity, but given their cumulative toxicity (4–8 months of treatment increases the likelihood of ototoxicity or nephrotoxicity) and the requirement for parenteral or intramuscular administration, they would rank lower than the previously described compounds.² Although experience is still limited, meropenem/clavulanate has demonstrated efficacy and is bactericidal. A recent phase 2 study²⁸ and observational studies²⁹ compel us to consider meropenem/clavulanate a core drug at the same level as the second-line aminoglycosides. Meropenem was found to be more active than imipenem in a recent study.³⁰

Ertapenem has been used successfully as switch therapy for cases treated at the hospital with meropenem who needed to be discharged, as it can be administered once daily intramuscularly.³¹

If the efficacy and safety profile of certain drugs like linezolid, bedaquiline, and delamanid is confirmed, they might move up the anti-TB drug hierarchy, as recently suggested.² At present bedaquiline and delamanid are recommended in XDR-TB or MDR-TB cases only, when no other options are available or tolerated.^{23,27}

Recognizing and promptly managing possible adverse events remains a priority, as recommended by the WHO,^{23,27} for the new drugs as well as for the other second-line drugs (e.g., thionamides, cycloserine, and *p*-aminosalicylic acid, among others).

Among the factors contributing to the programmatic success of a given drug, the availability of the drug and of the laboratory tests to confirm susceptibility or resistance, as well as the cost-effectiveness profile, is highlighted.

Establishing a hierarchy of second-line anti-TB drugs is difficult, as we try to evaluate the effect of a single component within a multidrug regimen where there is a lot of background noise.

3. Conclusions

In conclusion, the WHO has recently provided an important and useful evidence-based new classification of anti-TB drugs, which is the present roadmap allowing clinicians to correctly design safer and more effective MDR- and XDR-TB treatment regimens. As more evidence becomes available, further changes are likely to occur, particularly with the new drugs and some of the previous group 5 drugs.

It is hoped that ongoing RCTs will soon provide the necessary information to further improve the clinical and programmatic approach to the management of MDR- and XDR-TB cases.

Acknowledgements

This article forms part of the activities of the ERS/ALAT SinTB project.

The authors alone are responsible for the views expressed in this article; they do not necessarily represent the decisions and policies of the authors' institutions.

Financial support: No funding was received by the authors.

Conflict of interest: No competing interests declared.

References

- World Health Organization . WHO treatment guidelines for drug-resistant tuberculosis 2016 update. WHO/HTM/TB 2016.04. Geneva: WHO; 2016.
- Caminero JA, Scardigli A. Classification of antituberculosis drugs: a new proposal based on the most recent evidence. *Eur Respir J* 2015;**46**:887–93.
- World Health Organization. Companion handbook to the WHO guidelines for the programmatic management of drug-resistant tuberculosis. WHO/HTM/TB/2014.11. Geneva: WHO; 2014.
- Caminero JA, Sotgiu G, Zumla A, Migliori GB. Best drug treatment for multidrug-resistant and extensively drug-resistant tuberculosis. *Lancet Infect Dis* 2010;**10**:621–9.
- ElMaraachli W, Slater M, Berrada ZL, Lin SY, Catanzaro A, Desmond E, et al. Predicting differential rifamycin resistance in clinical *Mycobacterium tuberculosis* isolates by specific *rpoB* mutations. *Int J Tuberc Lung Dis* 2015;**19**:1222–6.
- Caminero JA. Guidelines for clinical and operational management of drug-resistant tuberculosis. Paris, France: International Union Against Tuberculosis and Lung Disease; 2013.
- Caminero JA, Matteelli A, Lange C. Treatment of TB. In: Lange C, Migliori GB, editors. *Tuberculosis (ERS Monograph)*. Sheffield: European Respiratory Society; 2012. p. 154–66.
- Ahuja SD, Ashkin D, Avendano M, Banerjee R, Bauer M, Bayona JN, et al. Multidrug resistant pulmonary tuberculosis treatment regimens and patient outcomes: an individual patient data meta-analysis of 9,153 patients. *PLoS Med* 2012;**9**:e1001300.
- Falzon D, Gandhi N, Migliori GB, Sotgiu G, Cox H, Holtz TH, et al. Resistance to fluoroquinolones and second-line injectable drugs: impact on MDR-TB outcomes. *Eur Respir J* 2013;**42**:156–68.
- Migliori GB, Sotgiu G, Gandhi NR, Falzon D, De Riemer K, Centis R, et al. ; Collaborative Group for Meta-Analysis of Individual Patient Data in MDR-TB. Drug resistance beyond extensively drug-resistant tuberculosis: individual patient data meta-analysis. *Eur Respir J* 2013;**42**:169–79.
- Guglielmetti L, Le Dú D, Veziris N, Caumes E, Marigot-Outtandy D, Yazdanpanah Y, et al. Is bedaquiline as effective as fluoroquinolones in the treatment of multidrug-resistant tuberculosis? *Eur Respir J* 2016;**48**:582–5.
- Sotgiu G, Centis R, D'Ambrosio L, Alffenaar JW, Anger HA, Caminero JA, et al. Efficacy, safety and tolerability of linezolid containing regimens in treating MDR-TB and XDR-TB: systematic review and meta-analysis. *Eur Respir J* 2012;**40**:1430–42.
- Sotgiu G, Pontali E, Migliori GB. Linezolid to treat MDR-/XDR-tuberculosis: available evidence and future scenarios. *Eur Respir J* 2015;**45**:25–9.
- Bolhuis MS, Tiberi S, Sotgiu G, De Lorenzo S, Kosterink JG, van der Werf TS, et al. Linezolid tolerability in multidrug-resistant tuberculosis: a retrospective study. *Eur Respir J* 2015;**46**:1205–7.
- Lewis JM, Hine P, Walker J, Khoo SH, Taegtmeier M, Squire SB, et al. First experience of effectiveness and safety of bedaquiline for 18 months within an optimised regimen for XDR-TB. *Eur Respir J* 2016;**47**:1581–4.
- Tadolini M, Lingsang RD, Tiberi S, Enwerem M, D'Ambrosio L, Sadutshang TD, et al. First case of extensively drug-resistant tuberculosis treated with both delamanid and bedaquiline. *Eur Respir J* 2016;**48**:935–8.
- Diacon AH, Dawson R, Von Groote-Bidlingmaier F, Symons G, Venter A, Donald PR, et al. Randomized dose-ranging study of the 14-day early bactericidal activity of bedaquiline (TMC207) in patients with sputum microscopy smear-positive pulmonary tuberculosis. *Antimicrob Agents Chemother* 2013;**57**:2199–203.
- Diacon AH, Pym A, Grobusch M, Patientia R, Rustomjee R, Page-Shipp L, et al. The diarylquinoline TMC207 for multidrug-resistant tuberculosis. *N Engl J Med* 2009;**360**:2397–405.
- Diacon AH, Pym A, Grobusch MP, de los Rios JM, Gotuzzo E, Vasilyeva I, et al. Multidrug-resistant tuberculosis and culture conversion with bedaquiline. *N Engl J Med* 2014;**371**:723–32.
- Pym AS, Diacon AH, Tang SJ, Conradie F, Danilovits M, Chuchottaworn C, et al. Bedaquiline in the treatment of multi- and extensively drug-resistant tuberculosis. *Eur Respir J* 2016;**47**:564–74.
- Tiberi S, De Lorenzo S, Centis R, Viggiani P, D'Ambrosio L, Migliori GB. Bedaquiline in MDR/XDR-TB cases: first experience on compassionate use. *Eur Respir J* 2014;**43**:289–92.
- van Halsema C, Humphreys S, Bonington A. Extensively drug-resistant tuberculosis: early access to bedaquiline for a UK patient. *Eur Respir J* 2014;**43**:292–4.
- World Health Organization. The use of bedaquiline in the treatment of multidrug-resistant tuberculosis: interim policy guidance. WHO/HTM/TB/2013.6. Geneva: WHO; 2013.
- Hartkoon RC, Uplekar S, Cole ST. Cross-resistance between clofazimine and bedaquiline through upregulation of MmpL5 in *Mycobacterium tuberculosis*. *Antimicrob Agents Chemother* 2014;**58**:2979–81.
- Skripconoka V, Danilovits M, Pehme L, Tomson T, Skenders G, Kummik T, et al. Delamanid improves outcomes and reduces mortality in multidrug-resistant tuberculosis. *Eur Respir J* 2013;**41**:1393–400.
- Wells CD, Gupta R, Hittel N, Geiter LJ. Long-term mortality assessment of multidrug-resistant tuberculosis patients treated with delamanid. *Eur Respir J* 2015;**45**:1498–501.
- World Health Organization. The use of delamanid in the treatment of multidrug-resistant tuberculosis. Interim policy guidance. WHO/HTM/TB/2014.23. Geneva: WHO; 2014.
- Diacon AH, van der Merwe L, Barnard M, von Groote-Bidlingmaier F, Lange C, García-Basteiro AL, et al. β -Lactams against tuberculosis—new trick for an old dog? *N Engl J Med* 2016;**375**:393–4.
- Sotgiu G, D'Ambrosio L, Centis R, Tiberi S, Esposito S, Dore S, et al. Carbapenems to treat multidrug and extensively drug-resistant tuberculosis: a systematic review. *Int J Mol Sci* 2016;**17**:373.
- Tiberi S, Sotgiu G, D'Ambrosio L, Centis R, Arbex MA, Alarcon AE, et al. Comparison of effectiveness and safety of imipenem/clavulanate- versus meropenem/clavulanate-containing regimens in the treatment of multidrug and extensively drug-resistant tuberculosis. *Eur Respir J* 2016;**47**:1758–66.
- Tiberi S, D'Ambrosio L, De Lorenzo S, Viggiani P, Centis R, Sotgiu G, et al. Ertapenem in the treatment of multidrug-resistant tuberculosis: first clinical experience. *Eur Respir J* 2016;**47**:333–6.